



ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

New Astronomy Reviews xxx (2004) xxx–xxx

**New Astronomy
Reviews**

www.elsevier.com/locate/newastrev

The ISM in nearby galaxies

Elias Brinks

INAOE, Puebla, Mexico

6 Abstract

7 The SKA will revolutionise the study of the principles underlying star formation (SF), resolving interstellar cloud
 8 complexes which are the birthplaces of stars and answering such questions as which are the sufficient and necessary
 9 conditions for SF to commence. Also, massive SF is intimately related to stellar death. The SKA will be able to study
 10 the structure of the ISM at 100 pc resolution out to distances of up to 20 Mpc and will quantify the impact the demise of
 11 massive stars has on their environment. Importantly, the SKA will probe the transition region between ISM and IGM,
 12 linking star formation and stellar death in the disks of galaxies to faint HI structures further afield, such as “anomalous
 13 gas” and (Compact) High Velocity Clouds. Lastly, the superb sensitivity of the SKA will result in some hundred back-
 14 ground sources per square degree against which HI absorption lines can be searched for, probing not only the relative
 15 importance of the different phases of the gas in galaxies but also the low density gas in the outskirts and between
 16 galaxies.

17 © 2004 Published by Elsevier B.V.

18

19 1. Understanding star formation

20 Star formation is arguably the single most
 21 important process to shape the Universe as we
 22 know it. The very first luminous objects, through
 23 their ionising radiation field, put an end to the
 24 “Dark Ages”. Star formation in the first structures
 25 that formed caused (proto-)galaxies to light up,
 26 creating the stunning, galaxy studded images in
 27 deep exposures revealed by HST. And it is star for-
 28 mation that enriches the interstellar and interga-
 29 lactic medium (ISM and IGM, respectively) with
 30 metals and which continues to change the balance

between the amount of matter within galaxies and
 the number of stars.

31
 32
 33 Despite the obvious importance of the star for-
 34 mation (SF) process, embarrassingly little is
 35 known about the sufficient and necessary condi-
 36 tion for SF to commence. To improve this situa-
 37 tion, high spatial and velocity resolution maps
 38 are required which resolve the individual cloud
 39 and cloud complexes of atomic hydrogen (HI)
 40 which will collapse to form stars. It is from these
 41 neutral cloud complexes that giant molecular
 42 clouds will condense. As their linear sizes are of
 43 order 100 pc, this requires an angular resolution of
 44 1" at a distance of 20 Mpc. In other words, in
 45 order to understand SF at a cosmological level,

E-mail address: ebrinks@inaoep.mx.

46 studies of nearby galaxies are indispensable and
 47 are our only way to address questions such as what
 48 triggers the onset of star formation: is it due to a
 49 local gravitational instability or does one need an
 50 external driver, for example an interaction, is it
 51 triggered by density waves or rather a bar instabil-
 52 ity? What is the role played by magnetic fields
 53 (Beck and Gaensler, this volume). Closely related
 54 to this is the question of how SF depends on gal-
 55 axy environment, galaxy type, galaxy kinematics
 56 and heavy element abundance of the ISM.

57 The SKA will be the only instrument capable of
 58 reaching the required resolution at sufficient sensi-
 59 tivity within an acceptable amount of observing
 60 time. Assuming that 50% of the total array collect-
 61 ing area will fall within a diameter of 6 km and
 62 75% within a diameter of 30 km, taking $A_{\text{eff}}/$
 63 $T = 20,000 \text{ m}^2 \text{ K}^{-1}$ between 0.5 and 1.5 GHz,
 64 and hence $A_{\text{eff}} = 2.5 \times 10^5 \text{ m}^2$ and taking as a
 65 maximum baseline 45 km in order to achieve a
 66 $\theta_{\text{syn}} = 1''$, a 12-h targeted observation at 10 kHz
 67 (2 km s^{-1}) velocity resolution will reach a 3σ limit
 68 on the surface brightnesses within a single channel
 69 of typically 10 K. A $\theta_{\text{syn}} = 1''$ corresponds to 0.25
 70 pc at the distance of the Magellanic Clouds (50
 71 kpc), 18 pc at the distance of the M81 group (3.6
 72 Mpc), and 50 pc at 10 Mpc.

73 There are over a thousand galaxies of all Hub-
 74 ble types within 10 Mpc. The Virgo cluster is in-
 75 cluded if extending the range to 20 Mpc at which
 76 distance the linear resolution is still an acceptable
 77 100 pc, comparable to the published studies in
 78 M31 and M33 (Brinks and Bajaja, 1986; Deul
 79 and den Hartog 1990).

80 In order to cover galaxies of all Hubble types
 81 over a wide range of surface brightness, metallicity
 82 and star formation activity, a sample of at least
 83 100 objects will be required, e.g., along the line
 84 of the *Spitzer Space Telescope* SINGS Legacy pro-
 85 ject (Kennicutt et al., 2003; Walter et al., 2004)
 86 which translates to a 1200-h project. A project of
 87 this nature would, of course, benefit from a large
 88 ($\sim 1^\circ$) instantaneous field of view as well as from
 89 multiple fields of view.

90 It is illustrative to show what can be done cur-
 91 rently and what the SKA will achieve. Fig. 1 is
 92 an HI peak brightness map of the LMC (Kim et
 93 al., 1999) at a resolution of $1'$ or 15 pc and repre-

94 sents one of the highest linear resolution maps of
 95 an object in the Local Group. The SKA will allow
 96 the resolution to be increased 60-fold in the case of
 97 the Magellanic Clouds, whereas the resolution cur-
 98 rently obtained in the Clouds will be achieved in
 99 galaxies within 4 Mpc. Moreover, with a resolu-
 100 tion of $1''$ one can do the same kind of studies in
 101 external galaxies out to 4 Mpc as what has been
 102 done with 100-m single dish telescopes such as
 103 the GBT or Effelsberg at the distance of the center
 104 of the Milky Way.

105 The view is that star formation occurs above a
 106 certain gas column density threshold (e.g., Kenni-
 107 cutt, 1989; Martin and Kennicutt, 2001). The SKA
 108 will be able to answer the question if there is a
 109 “universal” star formation threshold, and to what
 110 extent this threshold is a function of galaxy type
 111 and of heavy element abundance. Studies of low
 112 metallicity systems, such as dwarf irregular (dIrr)
 113 galaxies, can be used to extrapolate to SF in the
 114 unenriched early universe. This study will benefit
 115 from synergies with other future instruments such
 116 as ALMA which will provide the necessary high
 117 resolution observations of the molecular gas, at
 118 comparable resolution.

119 Integrated HI surface density maps will be com-
 120 bined with velocity dispersion maps and the derived
 121 rotation curves to calculate spatially resolved
 122 “Toomre- Q ” parameter maps for each galaxy
 123 (the Q parameter is a measure for the local gravita-
 124 tional balance; Toomre 1964). An example of what
 125 will be possible is presented in Fig. 2 for the galaxy
 126 NGC 6822 (de Blok and Walter, 2000). Using these
 127 kind of maps the question can be addressed what
 128 the importance is of local (disk or cloud instability)
 129 versus global effects (spiral density waves, tidal
 130 forces, magnetic fields) in triggering SF.

2. The violent interstellar medium 131

132 Star birth is intimately related to star death.
 133 The most massive stars (with masses larger than
 134 $8 M_\odot$) explode as type II supernovae and deposit
 135 of order 10^{51} erg of kinetic energy in their immedi-
 136 ate surrounding creating coronal gas filled, over-
 137 pressured bubbles which expand into the
 138 surrounding ISM. These bubbles show up as 138




Fig. 1. Peak HI surface brightness map by Kim et al. (1999) in green with overlaid an H α image (with the continuum subtracted) of the LMC in red.

139 empty regions in HI (HI bubbles or superbubbles,
140 depending on their size). A spectacular example is
141 again the LMC, an image of which is reproduced
142 in Fig. 3 (Kim et al., 2003). The expanding,
143 shocked rims where subsequent star formation is
144 likely to occur, are known as giant or supergiant
145 shells. The superbubbles and supergiant shells are
146 thought to be due to the combined effect of multi-
147 ple SNe occurring within a young stellar associa-
148 tion (i.e., within a short time span and within a
149 limited volume). The accumulated energy is suffi-
150 cient to blow superbubbles which grow larger than
151 the thickness of the disk of a typical spiral galaxy,
152 launching the expanding shell and enriched, hot
153 gas into the halo.

154 It is this process which helps maintain the veloc-
155 ity dispersion of the gas at its canonical value of
156 around 6–7 km s⁻¹ (at least at scales of several
157 hundred pc) and which has been named as a pos-

sible explanation for the High Velocity Clouds or
158 HVCs (Galactic fountain model; Shapiro and
159 Fields, 1976; Bregman, 1980). The velocity disper-
160 sion of the gas largely defines the thickness of the
161 gas disks in gas rich spiral and dwarf galaxies. It
162 has been shown that especially in dwarf irregular
163 galaxies (dIrr) the disks are thicker in relative
164 and absolute sense (Brinks et al., 2002), increasing
165 the probability for lines of sight toward high red-
166 shift objects to intersect these halos, giving rise
167 to DLy α lines. Similarly, the hot gas hurled into
168 the halo will eventually cool and condense, aug-
169 menting effectively the cross section of a spiral gal-
170 axy and providing an explanation for DLy α and
171 associated metal lines. Importantly, dIrr galaxies,
172 which are thought to be particularly abundant at
173 large lookback times in current bottom up scenar-
174 ios, might not be able to hold on to the expanding
175 material leading to an early enrichment of the
176




Fig. 2. HI surface brightness map based on ATCA observations of the dwarf irregular galaxy NGC 6822 (de Blok and Walter, 2000) shown in the top left panel. The high resolution rotation curve is shown in the top middle and is combined with the surface brightness and velocity dispersion maps to produce a “Toomre Q ” map (bottom left) at linear scales of 90 pc. Dark regions correspond to low Q , implying that the gas there is gravitationally unstable and will turn molecular, fragment and form stars. An H α map is shown in the lower right. The SKA will allow the determination of the “Toomre Q ” locally and investigate its behavior as a function of environment and galaxy type.

177 IGM (Mac Low and Ferrara, 1999; Ferrara and
178 Tolstoy, 2000).

179 The SKA will have a huge impact on this field
180 as it will have the sensitivity and resolution to
181 map extremely low column density gas around
182 nearby galaxies, exploring the origin of HVCs
183 and linking the existence of gas at large distances
184 from a galaxy with recent star formation. The fact
185 that this will come within reach of the SKA is due,
186 in part, to its high surface brightness sensitivity.
187 Opting for a synthesised beam of, say 45" this cor-
188 responds to 800 pc at 3.6 Mpc and 2.5 kpc at 10
189 Mpc; using a channel width of 10 km s⁻¹, a 24-h
190 observation will reach typical column densities of
191 4×10^{17} cm⁻² which reaches well below the limit

192 where HI is supposedly ionised by the meta-galactic
193 radiation field. If pockets of gas exist which
194 were somewhat denser and more resistant to ionisation,
195 a 1 h observation with the same beam and
196 velocity resolution corresponds to mass limits of
197 $400 M_{\odot}$ at 1 Mpc or $4 \times 10^4 M_{\odot}$ at 10 Mpc.

198 At these sensitivities (Compact) High Velocity
199 Clouds around external galaxies come within easy
200 reach (de Heij et al., 2002; Thilker et al., 2000) as
201 will be the so-called “anomalous” gas which has
202 been found in very deep maps of a few individual
203 galaxies (Schulman et al., 1996, 1997; Fraternali et
204 al., 2002). As mentioned, SKA observations will
205 have the sensitivity and resolution to investigate
206 the link between SF activity (and the aftermath




Fig. 3. Peak HI surface brightness map for the LMC (Kim et al., 2003). The gray scale intensity range is 0–136.7 K. This map is sensitive to the small HI clouds with the highest opacity along the line of sight. In such regions, the brightness temperature will approach the spin temperature of the HI. The image is similar to the ATCA-only image of Kim et al. (1998) and emphasizes the filamentary, bubbly, and flocculent structure of the ISM in the LMC. The locations of supergiant HI shells are indicated.

207 of stellar death) in the disks of galaxies and mater- 217
 208 ial in the halo, verifying the validity of the galactic 218
 209 fountain model. Moreover, the SKA will be able 219
 210 to map this low density gas and see if there is 220
 211 any link with gas-rich satellite galaxies orbiting 221
 212 the bigger galaxy and eventually being dragged to- 222
 213 wards it through dynamical friction. 223

214 3. HI in absorption – going to the limit

215 The SKA will revolutionise HI absorption stud- 228
 216 ies. For HI absorption, the sensitivity is ultimately 229

determined by the flux density of the background 217
 source (or in the case of the source being extended, 218
 its surface brightness) against which absorption is 219
 detected. Assuming a 12-h integration at 1 km s^{-1} 220
 velocity resolution (cold HI clouds will have nar- 221
 row line widths) one can reach an rms noise of 222
 $15 \mu\text{Jy}$ per channel. The SKA will thus be able to 223
 detect a 10% absorption feature ($\tau \approx 0.1$) against 224
 a 1 mJy background source at better than 6σ signal 225
 to noise (or a column density of $2 \times 10^{19} \text{ cm}^{-2}$ for 226
 $T_{\text{sp}} = 100 \text{ K}$ spin temperature gas). 227

The expected source density of sources brighter 228
 than 1 mJy, based on source counts at 20 cm wave- 229

length is $\sim 3 \times 10^5$ sterad⁻¹ (Hopkins et al., 1998). There will therefore be of order 100 background sources bright enough for absorption line studies behind M 33, some 500 sources in the field of M 31, and over 3000 behind the LMC. The best one can do currently with instruments like the VLA in the nearest galaxies like M 31 and M 33 is measuring absorption towards a dozen background sources (Braun and Walterbos, 1992; Dickey and Brinks, 1993). HI absorption measurements against background sources seen through the disks of gas rich galaxies can be used to extract crucial information regarding the spin temperature of the gas, T_{sp} , and the fraction of cool versus warm gas filling the ISM, both globally and as a function of position in a galaxy. This can only be done with the SKA.

HI absorption against extended background sources will reveal fine structure in the ISM, the limit of which is set only by the brightness temperature distribution (angular size) of the background source. Note that for these type of observations one can exploit the full resolution of the SKA. These observations can be linked to efforts by several groups (e.g., Gazol et al., 2001) to gain a theoretical understanding of the 3D and temperature structure of the ISM, involving turbulence. Most studies have been focussing on the Milky Way, understandably. With the SKA similar studies can be extended to other galaxies, notably the LMC/SMC and larger galaxies in the Nearby Universe.

4. Concluding remarks

The SKA will have a profound impact on HI studies of the Nearby Universe. For the same observing time, the SKA will provide maps at an order of magnitude higher angular resolution, twice the velocity resolution and a factor of perhaps two in improved sensitivity. SKA maps of nearby galaxies will provide the benchmark against which high redshift studies will be evaluated. The SKA will tackle such fundamental questions as what triggers star formation at the scale of neutral atomic clouds (~ 100 pc) and what is the

relation between violent star formation, and the death of massive stars, and the neutral gas which is seen at low column densities in the haloes of spiral galaxies. Is this gas due to outflow or are we rather seeing infall of gas, either stripped material from satellite galaxies which are in the process of merging, or primordial material which is still accreting. Lastly, HI absorption line studies will finally reveal what fraction of the ISM is in the form of cool ($T_{\text{sp}} = 80$ K) versus warm ($T_{\text{sp}} = 8000$ K) gas, addressing the validity of the fundamental and possibly erroneous assumption of the HI gas being optically thin.

References

- Braun, R., Walterbos, R.A.M., 1992. *ApJ* 386, 120.
 Bregman, J.N., 1980. *ApJ* 236, 577.
 Brinks, E., Bajaja, E., 1986. *A&A* 169, 14.
 Brinks, E., Walter, F., Ott, J., 2002. In: *Disks of Galaxies: Kinematics, Dynamics and Perturbations*, Athanassoula, E., Bosma, A., Mújica, R. (Eds.), ASP Conference Proceedings 275, p. 57.
 de Blok, W.J.G., Walter, F., 2000. *ApJ* 537, L95.
 de Heij, Braun, R., Burton, W.B., 2002. *A&A* 392, 417.
 Dickey, J.M., Brinks, E., 1993. *ApJ* 405, 153.
 Ferrara, A., Tolstoy, E., 2000. *MNRAS* 313, 291.
 Fraternali, F., van Moorsel, G., Sancisi, R., Oosterloo, T., *AJ* 123, 3124.
 Gazol, A., Vázquez-Semadeni, E., Sánchez-Salcedo, F.J., Scalo, J., 2001. *ApJ* 557, L121.
 Hopkins, A.M., Mobasher, B., Cram, L., Rowan-Robinson, M., 1998. *MNRAS* 296, 839.
 Kennicutt Jr., R.C., 1989. *ApJ* 344, 685.
 Kennicutt Jr., R.C., et al, 2003. *PASP* 115, 928.
 Kim, S., Staveley-Smith, L., Dopita, M.A., Sault, R.J., Freeman, K.C., Lee, Y., Chu, Y.-H., 2003. *ApJS* 148, 473.
 Kim, S., Dopita, M.A., Staveley-Smith, L., Bessell, M.S., 1999. *AJ* 118, 2797.
 Mac Low, M.-M., Ferrara, A., 1999. *ApJ* 513, 142.
 Martin, C.L., Kennicutt, R.C., 2001. *ApJ* 555, 301.
 Schulman, E., Brinks, E., Bregman, J.N., Roberts, M.S., 1997. *AJ* 113, 1559.
 Schulman, E., Bregman, J.N., Brinks, E., Roberts, M.S., 1996. *AJ* 112, 960.
 Shapiro, P.R., Fields, G.B., 1976. *ApJ* 205, 762.
 Thilker, D.A., Braun, R., Walterbos, R.A.M., Corbelli, E., Lockman, F.J., Murphy, E., Maddalena, R., 2000. *ApJ* 601, L39.
 Walter, F., Brinks, E., de Blok, W.J.G., Thornley, M., Kennicutt, R., 2004. *BAAS* 204 (submitted).

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323