

L183 (L134N) Revisited

III. The gas depletion^{*}

L. Pagani¹, J.-R. Pardo², A.J. Apponi³, A. Bacmann⁴, and S. Cabrit¹

¹ LERMA & UMR 8112 du CNRS, Observatoire de Paris, 61, Av. de l'Observatoire F-75014 Paris
e-mail: laurent.pagani@obspm.fr

² Instituto de Estructura de la Materia, Dpto de Física Molecular CSIC, Madrid, Spain

³ University of Arizona, Steward Observatory, 933 N. Cherry Ave., Tucson, AZ 85721

⁴ Observatoire de Bordeaux, BP 89, 33270 Floirac, France

Received 08/04/2004; Accepted 30/07/2004

Abstract. We present a detailed study of the gas depletion in L183 (= L134N) for a set of important species, namely, CO, CS, SO, N₂H⁺ and NH₃. We show that all these species are depleted at some level. This level seems to depend mostly on a density threshold rather than on dust opacity. Therefore UV shielding would not be a main factor in the triggering of depletion. Our data suggest that CO, CS and SO depletion happen at densities of $\sim 3 \times 10^4 \text{ cm}^{-3}$, while N₂H⁺ and NH₃ seem to deplete at densities close to 10^6 cm^{-3} . The latter result is consistent with the Bergin & Langer (1997) polar (H₂O) ice case but not with the more recent models of Aikawa et al. (2003). CS depletion occurs much below its (J:2–1) critical density, ($7 \times 10^5 \text{ cm}^{-3}$) and therefore makes this species unsuitable to study the density structure of many dark cloud cores.

Key words. ISM: abundances – ISM: molecules – ISM: dust, extinction – ISM: Structure – ISM: individual: L134N – ISM: individual: L183

1. Introduction

In the very cold, shielded interior of dark molecular clouds, most molecules except H-only species and possibly N-carriers are expected to stick to grains and thus disappear from the gas phase. This has been anticipated as early as a few years after the discovery of CO. Because CO was predicted to deplete onto grains at temperatures below 20 K, it was not clear why it could still be seen in 10 K clouds. Yet, no clear trace of depletion could be found for any species until facilities for observing and measuring cold dust became available. In the mid-nineties, bolometer arrays (MAMBO at 1.2 mm, SCUBA at 850 and 450 μm) have allowed astronomers to detect dense cold dust cores which had escaped all searches so far because C¹⁸O, which was supposed to trace gas column-density in cold molecular clouds and thus locate cores via their high mass content, was in fact depleted inside these quiescent cores. Cores were thus found from dust emission eventually and more recently via near or mid infrared dust absorption. As

a direct consequence, CO (and other molecules) depletion was eventually revealed. While depletion itself is no longer questioned, the physical conditions it requires for different species are not yet well-established. Polar molecules such as H₂O freeze onto grains at temperatures below ~ 100 K and apolar molecules (like CO) below ~ 20 K, but N-bearing species, such as NH₃ and N₂H⁺ do not seem to experience this as they are seen deep inside the cores and could even act as surrogate dust tracers. It is thus necessary to track depletion in as many cold clouds as possible to shed light on the details of the depletion mechanism.

A second reason to study depletion is linked to astrochemistry. Without taking depletion effects into account, all species are considered to be spatially coexistent when they are not (N₂H⁺ cannot coexist with CO e.g., Bergin & Langer 1997, Caselli 2002) and therefore relative abundances are not correctly evaluated. This is one of the reasons why previous abundance estimates made towards L183 by Swade (1989b) and by Dickens et al. (2000) should be considered very cautiously before checking astrochemical models against them. We have therefore undertaken to revisit this source in order to establish it as a true reference for chemical models.

Depletion was revealed in L183 for the first time by Pagani et al. (2002, hereafter Paper I) and independently confirmed by Juvela et al. (2002). However, in both works the depletion estimates were done by comparison with the ISOPHOT 100 and 200 μm data which only probe a thin

Send offprint requests to: L. Pagani

^{*} Based on observations made with the CFHT, the Iram 30-m and the ARO 12-m (formerly NRAO 12-m).

Table 1. Observational parameters and sensitivity

<i>Molecule</i>	<i>Transition</i>	<i>Frequency</i> (MHz)	<i>Beamsize</i> (arcsec)	<i>rms</i> ^a (K)
$C^{18}O$	$J:1-0$	109782.163	57 _{KP}	0.11
$C^{18}O^b$	$J:1-0$	109782.163	23 _I	0.059
$C^{18}O$	$J:2-1$	219560.327	28 _{KP}	0.26 ^e
$C^{18}O$	$J:2-1$	219560.327	12 _I	0.12 ^e
$C^{17}O$	$J:1-0$	112359.277	56 _{KP}	0.034
$C^{32}S$	$J:2-1$	97980.950 ^c	64 _{KP}	0.088
^{32}SO	$J_N:3_2-2_1$	99299.879	63 _{KP}	0.12
N_2H^+	$J:1-0$	93173.384 ^d	67 _{KP}	0.14
N_2H^+	$J:1-0$	93173.384 ^d	27 _I	0.037
N_2D^+	$J:2-1$	154217.096 ^d	16 _I	0.050
N_2D^+	$J:3-2$	231321.757 ^d	12 _I	0.097
$NH_2D_{para}^b$	$J_{KK'}:1_{11}-1_{01}$	110153.587 ^d	23 _I	0.054

^a median value over the map

^b data partly from Tiné et al. (2000)

^c Frequency is taken from Pagani et al. 2001

^d Hyperfine components

^e Small map towards Peak E only

_I IRAM 30-m telescope

_{KP} NRAO/ARO KP 12-m

outer layer (0–15 A_V) of cold (12–17 K) dust. Therefore the depletion along the line-of-sight was evaluated to a factor of 1.5–2 only. As we showed in a subsequent paper (Pagani et al. 2004, hereafter Paper II), these early estimates are not correct because they ignore the bulk of the dust, which is at $T < 10$ K and does not emit at 200 μm . Therefore, depletion has been strongly underevaluated in these earlier works. We have now obtained (Paper II) a composite Mid- and Near- Infrared dust absorption map which traces all the dust column density, together with a map of the dust emission at 1.2 mm, and therefore we can work out more precisely the depletion effects in this source.

In this paper, we will present in Sect. 2 the molecular line observations we have made, analyze them in Sect. 3 and discuss the depletion effects we find by comparison with our dust map from Paper II in Sect. 4. In Sect. 5, we will draw our conclusions.

2. Observations

We have observed four species, namely CO, CS, SO and N_2H^+ , and for each species at least two isotopomers and for each isotopomer at least 2 transitions (except for the N_2H^+ isotopomer for which we have only a few $J:3-2$ observations with low signal-to-noise ratio). For the goals of the present work, we concentrate on a small subset of these observations (Table 1) complemented with observations from Tiné et al. (2000). The remaining observations will be published in subsequent papers. The reference position of the source in this study is : 15h54m06.6s, $-2^\circ52'19''$ (J2000) which is the L134N reference position given by e.g. Swade (1989a) and the LSR velocity is +2.5

km s^{-1} . Note however that the SIMBAD position for this source is 15h54m09.2s, $-2^\circ51'39''$ (J2000) and that a few other reference positions are also found in the literature. We would suggest using the central position of the dust peak from now on as the reference in further work: 15h54m08.5s, $-2^\circ52'43''$ (J2000).

The subset of observations presented here have been carried out with either the Kitt Peak 12-m (formerly owned by NRAO, now operated by the University of Arizona, hereafter KP 12-m) or the IRAM 30-m.

2.1. Kitt Peak 12-m data

Observations started in December 1999 at Kitt Peak and spanned a seven month period. Complementary observations were obtained in January 2001. The Kitt Peak 12-m is equipped with two-polarization receivers at 1.3, 2 and 3 mm with filter banks and a versatile autocorrelator in parallel as backends. We used 100 kHz filters all the time and the 32K-channel autocorrelator. Pointing (and focus) were checked on planets when available or 3C273 and 3C279 and found to be stable within 5". Pointing was checked once for short observing periods and twice for long periods (up to 7 hours). Except for N_2H^+ , all observations were conducted in frequency switch mode. The N_2H^+ ($J:1-0$) line spans almost 20 km s^{-1} and is thus better observed in position switch mode. The off position was taken 10' northwest of the source where no $C^{18}O$ emission is seen. The reference position was observed every 30 minutes or so to check for gain variations. For the weak $C^{17}O$ line, we made long integrations, and gain was checked only once per session. Gain variations towards the reference position would have been undetectable for such low SNR spectra. Autocorrelator sampling was set to 6 kHz (12 kHz resolution) which represents 30 to 36 m s^{-1} velocity resolution depending on the frequency. Data were Hanning smoothed afterwards to velocity resolution up to 0.1 or 0.2 km s^{-1} depending on the SNR needed. Whenever possible, the flux scale was checked on NGC 2264 (for SO lines, e.g.) or IRC+10216 at the beginning of the observations. Mapping was done with 1' steps except in some critical places (such as the dust core) where 30" step sampling was preferred. We mapped most of the lines on an area 15' \times 15', large enough to cover the entire L183 core (but the cloud connects with L134 southward). In a few cases, we stopped the mapping before, when the line became too weak (N_2H^+ in most directions and ^{32}SO northward). The single side-band system temperature ($T_{\text{sys}}(\text{SSB})$) was in the range 100–200 K.

2.2. IRAM 30-m data

IRAM 30-m data were collected in April, May, December 2003 and May 2004. The IRAM 30-m is equipped with two polarization receivers at 1.1, 1.3, 2 and 3 mm. It has filter banks and a versatile autocorrelator. The system allows to observe up to 4 different lines simultaneously. We




Fig. 1. Comparison of $C^{18}O$ (J:1-0) (left) and (J:2-1) (right) spectra (towards peak E) from the Kitt Peak (KP) 12-m and IRAM 30-m telescopes after smoothing the latter to the KP 12-m resolution ($1'$ for the (J:1-0) and $30''$ for the (J:2-1) transitions). Coordinates are offsets in arcmin from reference position. X axis (velocity) scale is from 0 to 5 km s^{-1} , and Y axis (T_r^*) scale is from -0.5 to 4.0 K (J:1-0) and from -0.3 to 2.2 K (J:2-1)

worked with a frequency resolution of 20 kHz at 2 and 3 mm , giving us a velocity resolution around 50 m s^{-1} at 3 mm and 35 m s^{-1} at 2 mm . At 1.3 mm , we worked with 40 kHz resolution to keep the same 50 m s^{-1} velocity resolution. Pointing and focus were checked every 90 minutes on close-by continuum sources. The observations have been performed in frequency switch mode. We observed a few positions in N_2H^+ and N_2D^+ in April and May 2003. T_{sys} (SSB) were 100 K at 3 mm , 300 K at 2 mm and 400 to 700 K at 1.3 mm . The reader is referred to Tiné et al. (2000) for the description of their observations. We performed further observations of the $N_2D^+(J:2-1)$ line in December 2003 to improve and enlarge Tiné's map to a size of $100'' \times 200''$. We also mapped the $N_2D^+(J:3-2)$ transition. For both transitions, T_{sys} were similar to what we obtained in Spring 2003. The 1.3 mm receiver we used was HERA (9 pixels spaced by $24''$) with the new autocorrelator, VESPA. We observed the $N_2D^+(J:2-1)$ line with 10 kHz resolution ($= 20 \text{ m s}^{-1}$) and the $N_2D^+(J:3-2)$ line with 40 kHz resolution ($= 50 \text{ m s}^{-1}$).

2.3. Calibration

The KP 12-m telescope forward scattering and spillover efficiency (η_{fss} , Kutner & Ulrich, 1981) has been measured to be 0.64 at all wavelengths. This efficiency allows to calibrate our data in the T_r^* scale if we suppose the beam coupling to the source to be 1. Though the exact definition of the T_r^* scale depends on what angular size we use to measure η_{fss} , its general acceptance (or its closest approximation, η_{moon}) applies to calibrate signals

from extended sources larger than the main beam, such as L183 (see Teyssier et al. 2002 for a recent discussion on this point). For the IRAM 30-m, the forward efficiency (η_f) is identical to η_{moon} which means that for extended sources such as L183, it is less erroneous to use η_f than to use the main beam efficiency, η_{MB} , which introduces too large a correction. In other words, for the IRAM 30-m, $T_a^* \approx T_r^*$.

Indeed, we find (Fig. 1) that for $C^{18}O$ (J:1-0) and (J:2-1), the IRAM 30-m telescope gives comparable results to the KP 12-m telescope after convolution at the KP 12-m resolution if we use the IRAM 30-m T_a^* scale and the KP 12-m T_r^* scale. This indicates that the calibration of both telescopes is consistent for extended emission tracers. If we had applied the η_{MB} correction for both telescopes then the IRAM 30-m (J:1-0) line would have been 23% stronger than the KP 12-m line while the KP 12-m (J:2-1) line would have been 26% stronger than the IRAM 30-m line, which would have lead to considerable discrepancies in the (J:2-1)/(J:1-0) line ratios.

3. Results

Because of its importance in this work, we redisplay first the dust map (Fig. 2) we obtained in Paper II with arrows marking both the places of the molecular peaks (C, N, S and W) described by Dickens et al. (2000) to which we have added a fifth molecular peak, E (see below), and the places of the nearby dust peaks which we have numbered from 1 to 5.




Fig. 2. Dust map from Paper II (in log grey scale). Coordinates are J2000 equatorial. Letters C, E, N, S, and W mark the molecular peaks while numbers 1 to 5 mark the corresponding dust peaks (see text). Dust peak #2 is not visible in this map but only in the dust emission map at 1.2 mm (see Paper II). Contours are from 5 to 50 by steps of 5 mag (black contours) and 60 to 120 by steps of 30 mag (white contours)




Fig. 3. $C^{18}O$ (J:1-0) integrated intensity contours (from the KP 12-m telescope) superimposed on the dust map from Paper II (in log grey scale). Coordinates are J2000 equatorial. $C^{18}O$ levels : 0.2 to 1.8 by 0.2 $K km s^{-1}$ (r.m.s. $\approx 35 mK km s^{-1}$). The white and black dashed contour represents the $A_V = 20$ mag level. Letters C, E, N, S, and W mark the molecular peaks (see text) and the KP 12-m beam size is displayed in the top right corner

3.1. Large-scale Kitt Peak maps

Figs 3–7 show $C^{18}O$, $C^{17}O$, $C^{32}S$, ^{32}SO , and N_2H^+ emission intensity superposed as contours on the dust map




Fig. 4. same as Fig. 3 for $C^{17}O$ (J:1-0). $C^{17}O$ levels : 0.1 to 0.7 by 0.1 $K km s^{-1}$ (r.m.s. $\approx 30 mK km s^{-1}$)




Fig. 5. same as Fig. 3 for $C^{32}S$ (J:2-1). $C^{32}S$ levels : 0.1 to 1.1 by 0.2 $K km s^{-1}$ (r.m.s. $\approx 45 mK km s^{-1}$)

image from Paper II (see Fig. 2). For the dust map, the $A_V = 20$ mag contour is emphasized.

For both $C^{18}O$ (Fig. 3) and $C^{17}O$ (Fig. 4) (J:1-0) lines, it is clear that none of the four molecular peaks (N, S, C, W) described by Dickens et al. (2000) is visible (except a very small $C^{18}O$ peak towards Peak C but with no counterpart in $C^{17}O$). To the east, there is a $C^{18}O/C^{17}O$ peak which coincides with our dust peak # 5 (Fig. 2 and Paper II) and which we label E to keep consistency with Dickens et al. (2000, this peak is outside their $C^{18}O$ map). Interestingly enough, this dust peak # 5 reaches an optical extinction of $A_V \approx 25$ mag. While $C^{18}O$ does show opacity effects (as mentioned in Paper I), it is not the case for $C^{17}O$ (the main hyperfine component is approximately ten times weaker than the $C^{18}O$ line). Both isotopomers




Fig. 6. same as Fig. 3 for ^{32}SO ($J_N:3_2-2_1$). ^{32}SO levels : 0.3 to 2.1 by 0.3 K km s^{-1} (r.m.s. $\approx 60 \text{ mK km s}^{-1}$)




Fig. 7. same as Fig. 3 for N_2H^+ ($J:1-0$). N_2H^+ levels : 0.4 to 2.8 by 0.4 K km s^{-1} (r.m.s. $\approx 75 \text{ mK km s}^{-1}$)

show the same features and thus the absence of the four N, S, C and W peaks is not due to opacity effects but is real.

The C^{32}S ($J:2-1$) line emission map (Fig. 5) shows no correlation with any of the five peaks but only a general increase towards the center of the cloud. However, the emission peak is situated north-west of peak W, noticeably outside of the dust cloud core and the line is obviously optically thick as its self-reversed profile indicates¹ and the question of peak hiding in the map arises again.

¹ Dickens et al. 2000 advocated the existence of two independent peaks instead of a self-absorption feature, due to the use of an erroneous CS frequency, see Pagani et al. (2001).

The ^{32}SO ($J_N:3_2-2_1$) line emission map (Fig. 6) shows a good correlation with two of the peaks (W and S) with a small extension towards peak C. It does not extend as far to the north as the CO and CS lines but is otherwise as extended as C^{18}O . However, the strongest SO peak is the W peak and not the C peak while the dust content is 4 times as high towards dust peak 1 than towards dust peak 3 (see Fig. 2).

Finally, the N_2H^+ ($J:1-0$) map (Fig. 7) is clearly different from the previous four as its extent is much more limited. It is rather well correlated with the dust content above $A_V = 20$ mag with two prominent peaks coincident with molecular peaks N and C and two extensions reaching peaks W and S. Peak E is conspicuously absent (a very weak line, 0.15 K peak intensity - a 4σ detection - has been detected though). It is remarkable then that the only $\text{C}^{18}\text{O}/\text{C}^{17}\text{O}$ peak coinciding with a dust peak is the only peak which hardly emits in N_2H^+ ($J:1-0$). Overall, CO and N_2H^+ maps are completely unrelated.

While all these differing features could have looked puzzling fifteen years ago when looking at Swade's maps (Swade, 1989a), we now know that the main reason behind these differences is the depletion of CO, CS and SO onto grains, as we will discuss below. Tafalla et al. (2002) recently reached the same conclusions towards other pre-stellar cores.

3.2. Small scale maps and cut

These maps concern data taken with the IRAM 30-m telescope with resolutions in the range $11-23''$.

Tiné et al. (2000) mapped a small region around the central molecular peak (peak C) with the IRAM 30-m which we redisplay in Fig. 8 along with the main core dust peak (Pagani et al. 2003, Paper II). The molecular data corresponds to C^{18}O ($J:1-0$), N_2D^+ ($J:2-1$) and ($J:3-2$) and para- NH_2D ($J_{KK'}:1_{11}-1_{01}$) integrated line emission. The N_2D^+ ($J:2-1$) map differs from the Tiné et al. (2000) original one because we display here our own observations instead of theirs. We also mapped the N_2D^+ ($J:3-2$) line in December 2003 (a line which they failed to detect). C^{18}O shows almost no variation and no correlation with the dust peak while the two deuterated species peak approximately at the central molecular peak (peak C)², with only a weak extension towards the dust peak.

To better reveal the core depletion, we have also plotted these N-bearing species integrated intensities in a north-south cut across the dust peak (Fig. 9). We have also included some N_2H^+ ($J:1-0$) data from the KP 12-m map and from sparse IRAM 30-m observations, not numerous enough to make a map. The intensity axis is arbitrary and is set to have approximately all species integrated intensities proportional to the dust emission on both sides of the dust peak. The dust peak clearly emerges outside the molecular peak.

² The low resolution of Dickens et al. 2000 data probably explain that their peak seems slightly displaced eastward




Fig. 8. a–e. IRAM 30-m maps of the molecular peak C and dust peak 1 (see paper II and Dickens et al. 2000). **a** Dust peak emission measured at 1.2 mm. Levels are 9 to 24 by 3 MJy/Sr. **b** $C^{18}O$ ($J:1-0$) integrated intensity. Levels are 1.2 to 1.8 by 0.2 K km s⁻¹. **c** NH_2D ($J_{KK'}:1_{11}-1_{01}$) integrated intensity. Levels are 0.15 to 1.05 by 0.15 K km s⁻¹. **d** N_2D^+ ($J:2-1$) integrated intensity. Levels are 0.2 to 1.4 by 0.2 K km s⁻¹. **e** N_2D^+ ($J:3-2$) integrated intensity. Levels are 0.05 to 0.35 by 0.05 K km s⁻¹. The horizontal dashed line indicates where the dust peak should appear in declination for the maps of the different species and the cross marks the Dickens et al. (2000) molecular peak C. The black spots in the bottom right corners indicate the different map resolutions of the IRAM 30-m (from 11'' to 23''). R.m.s. is ≈ 24 mK km s⁻¹ for maps b and c, 40 mK km s⁻¹ for map d and 28 mK km s⁻¹ for map e

4. Depletion effects

Two difficulties arise in the search for depletion effects. The first one is to have a correct estimate of the dust content. For example, Willacy et al (1998) have used ISOPHOT 100 and 200 μ m data to estimate the dust content in the very cold source L1498. While the $C^{18}O$ depletion is unquestionable in that source, the dust is probably largely underestimated (as noted by the authors themselves) because the 200 μ m emission does not trace dust below 10 K as we also show in Paper II. As mentioned above, we (Paper I) and Juvela et al. (2002) met the same problem. The second difficulty is to have a correct estimate of the species column density. For example, in cold clouds, with high gas column densities, $C^{18}O$ can become optically thick. In fact, if it were not for depletion effects, optically thick $C^{18}O$ would be very wide-spread in dark clouds with optical extinction above $A_V \approx 20$ mag. One solution is to use $C^{17}O$ instead but the line is weak and difficult to map on large areas. Another solution is to map the $C^{18}O$ ($J:2-1$) and ($J:3-2$) transitions and run detailed radiative transfer models. It is a time-consuming task too and the result is not straightforward as it implies some interpretation of the data prior to diagnosing depletion effects. The question thus often arises to know whether the $C^{18}O$ intensity flattening commonly observed towards the cores of cold dark clouds is really due to depletion or instead to radiative transfer effects such as optically thick gas or extremely cold (5–6 K) gas which could induce the same behaviour. Tafalla et al. (2002) have shown that depletion is the only possible explanation for the five starless

cores they have studied and most published cases usually show equivalent $C^{17}O$ behavior which rules out opacity effects. Low temperatures are usually thought difficult to justify from NH_3 and CO measurements. Here we are going to discuss depletion effects for CO, CS and SO on the large scale and then the depletion effects we see with nitrogen-bearing species (N_2H^+ , N_2D^+ , and NH_2D).

4.1. CO depletion

If we plot the pixel-to-pixel comparison of the $C^{18}O$ integrated intensity versus the dust optical extinction and combine the two sets of $C^{18}O$ data (KP 12-m and IRAM 30-m), we obtain the very clear-cut picture of depletion of Fig. 10. The A_V data (Paper II) have been smoothed to each telescope resolution and thus the peak opacity is only ~ 100 mag now for a 23'' beam and ~ 50 mag for a 60'' beam. The turn-over is abrupt and occurs at $A_V \approx 20$ mag. This is a factor of 2–3 higher than the value found by Bergin et al. (2002) in B68 where $C^{18}O$ saturation starts at $A_V \approx 7$ mag and becomes flat at $A_V \approx 10$ mag, or by Kramer et al. (1999) in IC 5146 and by Alves et al. (1999) in L977 where depletion is found to start at $A_V \approx 10$ mag for both sources. Thus except for peak E (or dust peak 5) all the region inside the $A_V = 20$ mag contour is depleted of CO isotopomers. The $C^{18}O$ depletion surface covers $\sim 25'^2$, a wide area indeed which explains the poor resemblance between the $C^{18}O$ map and maps of several other species in this source. A possible marginal increase of the $C^{18}O$ integrated intensity is still visible beyond the




Fig. 9. North-south cut through the dust peak (R.A. offset = +30''). The histogram displays the 1.2 mm dust emission while the sets of dots trace N-bearing species through the peak. The vertical scales are all arbitrary. They are set to have the species integrated intensities proportional to the dust emission on the peak sides. Except when specified, all data come from the IRAM 30-m. For NH_2D and N_2D^+ ($J:2-1$), we have also plotted the sum of all the optically thin hyperfine components (dotted lines and crosses) while for N_2H^+ ($J:1-0$), we have plotted the only hyperfine component ($(J_{F_1F_2}:1_{11}-0_{01})$, dashed line and crosses) which remains optically thin ($\tau \leq 1$) in this cut. Error bars are 1σ errors on the integrated intensities

depletion turnover. A linear fit is not physical because the depletion probably increases going deeper into the core but the slope gives a good lower limit to the depletion we observe here. We have fitted the Kitt Peak data for $A_V < 20$ mag on the one hand and the whole IRAM data on the other hand (Fig.10). We find a slope of 87 and




Fig. 10. C^{18}O ($J:1-0$) integrated intensity pixels cross-correlated with the A_V dust map pixels from Paper II. A_V map resolution is degraded to 1' for KP 12-m data and to 23'' for IRAM 30-m data. The slope variation indicates that the depletion reaches at least a factor 43 in volume

2.0 $\text{mK km s}^{-1} \text{mag}^{-1}$ respectively. The decline in abundance from the region below to the region above $A_V \approx 20$ mag is thus larger than ~ 43 on average (and the line of sight depletion is about 5 for the peak opacity). Below $A_V \approx 20$ mag, the C^{18}O abundance is $\sim 1.1 \times 10^{-7}$, though the detailed Monte-Carlo analysis performed for two positions (Paper I) and which yields higher column densities than the simpler LVG analysis indicates a C^{18}O abundance close to 1.4×10^{-7} , quite comparable to the value given by Frerking et al. (1982).

As discussed in Sect. 3, the comparison between C^{18}O and C^{17}O maps shows identical features between the two isotopomers and thus the opacity of the C^{18}O ($J:1-0$) is relatively small and cannot explain the absence of molecular peaks towards the main dust peak. Low temperatures are not a better explanation since, on the one hand, 40'' away from the dust peak, i.e. towards the L134N reference position, we have estimated the C^{18}O total column density to be $2.7 \times 10^{15} \text{cm}^{-2}$ (Paper I), and, on the other hand, the line intensities of the C^{18}O transitions change very little between the two positions, and therefore the C^{18}O column density should be approximately the same towards the dust peak. From the dust peak estimated A_V (~ 100 mag for 23'' resolution) and using $X(\text{C}^{18}\text{O}) = 1.4 \times 10^{-7}$, we would have expected a total C^{18}O column density of $\sim 1.4 \times 10^{16} \text{cm}^{-2}$ in that direction if there were no depletion. A mean temperature of ~ 6 K along the line of sight would be compatible with the observations but is unrealistic taking into account the envelope higher temperature (10-12 K). We would be forced to put the core close to 3 K to prevent excess C^{18}O emission. On top of that, the ($J:1-0$) line opacity would be ~ 10 , which is not compatible with the C^{17}O measurements. We would also have obtained 1.7 K for the main C^{17}O component instead of the observed 0.4 K. Depletion is indeed the best explanation we presently know of.




Fig. 11. $C^{32}S$ (J:2–1) and $C^{18}O$ (J:1–0) integrated intensity cross-correlation. KP 12-m data only. Square symbols represent the average of the $C^{32}S$ values per 0.2 K km s^{-1} $C^{18}O$ intervals. Error bars represent the 1σ dispersion. The straight line goes through the origin to the 9th average value, its slope is 0.48




Fig. 13. ^{32}SO ($J_N:3_2-2_1$) and $C^{18}O$ (J:1–0) integrated intensity cross-correlation. KP 12-m data only. Square symbols represent the average of the ^{32}SO values per 0.2 K km s^{-1} $C^{18}O$ intervals. Error bars represent the 1σ dispersion. The straight line goes through the origin to the 9th average value, its slope is 0.78




Fig. 12. $C^{32}S$ (J:2–1) integrated intensity (KP 12-m data) cross-correlated with A_V data degraded to $1'$ resolution

4.2. CS depletion

Contrarily to $C^{18}O$, the CS line is optically thick almost everywhere and the question is whether the opacity hides the dust peaks or if depletion prevents CS from revealing them. In fact, opacity is not the good reason : as Penzias (1975) has shown (see also Linke et al. 1977), if a line is optically thick and subthermally excited, then its intensity is merely proportional to its column density (this is illustrated in Appendix A). However, this is true only for a uniform gas slab. Here, a dilution of the peak emission in the surrounding gas is still to be expected and Penzias' prediction is only to be considered on a large scale. However, the CS peak emission is noticeably asymmetrical compared to the cloud core (Fig. 5), and a preliminary Monte-Carlo analysis has confirmed that the observed CS emission can be explained with densities lower than the critical density (Paper I). Therefore, we think that despite its optical depth, the $C^{32}S$ (J:2–1) map is not hiding the peaks (a

limited preliminary optically thin $C^{34}S$ (J:2–1) map confirms this point).

Though the CS map (Fig. 5) is somewhat different from the $C^{18}O$ map (Fig. 3), apart from their similar extent, their intensities are mostly proportional (Fig. 11). The proportionality factor is 0.48 and the reason is that $C^{18}O$ is easier to thermalize than CS. As mentioned above, this is due to the fact that CS (J:2–1) is subthermally excited and optically thick and thus its intensity is proportional to its column density. On the contrary, the $C^{18}O$ integrated intensity is also proportional to its column density because the (J:1–0) transition is thermalized and optically thin in most of the considered range. Therefore it is not surprising to find that CS has the same depletion behaviour as $C^{18}O$ (Fig. 12). CS depletion has been observed before (see e.g. Tafalla et al. 2002). A preliminary radiative transfer analysis (Paper I) has shown that the $C^{32}S$ and $C^{34}S$ transitions (J:1–0), (J:2–1) and (J:3–2) towards the L134N reference position can be explained by densities $\leq 3 \times 10^4 \text{ cm}^{-3}$, more than an order of magnitude below the CS (J:2–1) critical density ($n_c = 7 \times 10^5 \text{ cm}^{-3}$). These findings are consistent and point to the fact that CS cannot be used as a density tracer at all in cold dark clouds because it is depleted much below its critical density. The sudden drop for $\langle I(C^{18}O) \rangle = 1.9 \text{ K km s}^{-1}$ average will be discussed in Sect. 4.4.

4.3. SO depletion

Comparison of SO versus $C^{18}O$ (Fig. 13) shows that SO hardly appears below a $C^{18}O$ intensity of 0.6 K km s^{-1} (equivalent to $A_V = 5 \text{ mag}$) and that there is no proportionality above that limit (again, the sudden drop for $\langle I(C^{18}O) \rangle = 1.9 \text{ K km s}^{-1}$ will be discussed in Sect. 4.4). Its emission strength correlates somewhat with the C, W and S peaks (Fig. 14), but is not, or only weakly, corre-




Fig. 15. **left** ^{32}SO ($J_N:3_2-2_1$) integrated intensity cross-correlated with C^{32}S ($J:2-1$) integrated intensity (KP 12-m data). Square symbols represent the average of the ^{32}SO values per 0.1 K km s^{-1} C^{32}S intervals. Error bars represent the 1σ dispersion. The straight line is the 1:1 slope. **right** ^{32}SO ($J_N:3_2-2_1$)/ C^{32}S ($J:2-1$) integrated intensity ratio cross-correlated with A_V data degraded to $1'$ resolution. Square symbols represent the average of the SO/CS ratio per 5 mag intervals, except the last symbol which is the average of all the 5 points above $A_V = 32.5$ mag




Fig. 14. ^{32}SO ($J_N:3_2-2_1$) peak intensity (filled black contours) with N_2H^+ ($J:1-0$) integrated intensity (grey contours) superimposed to it. Coordinates are offset from the reference position. ^{32}SO levels : 0.3 to 2.4 by 0.3 K. N_2H^+ levels : 0.4 to 2.8 by 0.4 K km s^{-1}

lated with peaks N and E which could mean that it comes closer to the C, W and S peaks than CS. Interestingly enough, Bergin & Langer (1997) find that after about 1 My of evolution, when C and consequently CS have already started to deplete, there is a strong enhancement of SO before it starts depleting too. This enhancement has an amplitude of almost two orders of magnitude and SO eventually becomes more abundant than CS, while the model starts with a CS/SO ratio of ~ 100 . Aikawa et al. (2003, their Figs 3 to 5) show a similar SO enhancement in some

of their depletion models as a function of core distance, though the effect does not reach the same amplitude; i.e. SO never becomes more abundant than CS but the increase of its column density as a function of radial proximity to the core center is clearly visible. We have some hints of this effect here but we can only do a preliminary analysis for the following reason : there are no SO collision coefficients yet at low temperatures ($< 50 \text{ K}$, Green 1994) to make a non-LTE analysis of its excitation and thus only a rotational diagram analysis is presently possible. However, due to opacity effects, as explained by Goldsmith & Langer (1999), and to low excitation temperature³, this analysis is difficult to perform and requires many lines. We have shown that without proper corrections, SO column density would be underestimated by 2 orders of magnitude using such a diagram (Paper I). Though we presently lack data to study in detail the CS/SO column density ratio in L183, we have evaluated the CS and SO column densities towards the reference position (Paper I) for which we have observed many SO transitions and could gather a few more data from the literature. We have found $N(\text{CS}) \approx 1.4 \times 10^{13} \text{ cm}^{-2}$ and $N(\text{SO}) \approx 2.5 \times 10^{14} \text{ cm}^{-2}$, or a CS/SO ratio ≈ 0.056 . In addition, CS is more extended than SO (Figs 5 & 6) implying that away from the depleting zones, CS is probably more abundant than SO. To check this point we have compared the SO and CS integrated intensities, and the ratio of SO ($J_N:3_2-2_1$) to CS ($J:2-1$) line integrated intensities with A_V (Fig. 15). Both lines are emitted in regions with hydrogen density below their critical density and both lines are optically thick, implying that they are both proportional to their column

³ As shown in Fig. 3 of Paper I, the cosmic background is no longer negligible if the excitation temperature is close to 2.73 K, i.e. $T_{ex} \leq 5 \text{ K}$, which is the case of most species with high dipolar momentum in the typical density range of dark clouds.




Fig. 17. N_2H^+ ($J:1-0$) spectrum taken towards the dust peak with the IRAM 30-m. The dashed curve is the CLASS hyperfine Gaussian fit result, the (1:3:5:7) LTE ratio of the hyperfine components is shown in the top right corner

density to a first approximation. There is a clear trend for SO to start at a comparable or slightly lower intensity than CS and, coming closer to the depleting core, to become more intense up to $A_V = 15$ mag. This obviously requires a more quantitative evaluation.

4.4. N-bearing species

It has been recognized that N-bearing species like NH_3 and N_2H^+ should deplete only at very high densities, $n > 10^6 \text{ cm}^{-3}$ (Bergin & Langer 1997), or even higher (e.g. $n \geq 3 \cdot 10^7 \text{ cm}^{-3}$ according to Aikawa et al. 2003). The reason given is the lower binding energy for Nitrogen with respect to Carbon or Oxygen. This ability to stay longer in the gas phase should make these species good mass tracers in the dark cloud cores. Furthermore deuterated N-bearing species should be even more efficient core tracers, deuteration being enhanced by the disappearance of CO (e.g. Caselli 2002, Bacmann et al. 2003). However some question has arisen as to whether these species should eventually also start depleting in the densest, coldest pre-stellar cores as reported by Bergin et al. (2002). We have thus compared our N_2H^+ ($J:1-0$), N_2D^+ ($J:2-1$) and ($J:3-2$) and the para- NH_2D ($J_{KK'}:1_{11}-1_{01}$) data of Tiné et al. (2000) to our dust map (Fig. 8).

4.4.1. N_2H^+

The N_2H^+/A_V comparison (Fig. 16a) shows that N_2H^+ appears above $A_V \approx 10$ mag only and also shows signs of depletion above $A_V \approx 50$ mag. The north-south cut across the main dust peak (Fig. 9) also suggests depletion (more correctly, the depletion concerns the parent species of N_2H^+ , N_2 , as the ion cannot stick to grains itself). However the line is strongly saturated as the spec-

trum taken towards the dust peak shows (Fig. 17) and it could be argued that the saturation above $A_V \approx 50$ mag in Fig. 16a comes from opacity effects. To check this point, we have studied the ($J_{F_1F_2}:1_{11}-0_{01}$) line and estimated its opacity using the CLASS HFS method⁴. We find total opacities (sum of the opacities of all the 7 hyperfine transitions of the $J:1-0$ line) in the range 10–30 mostly, yielding ($J_{F_1F_2}:1_{11}-0_{01}$) opacities in the range 0.4–1.1. This is the weakest and the only optically thin hyperfine transition for this ($J:1-0$) line and this range of total opacities. We have also compared its integrated intensity to A_V (Figs 9 & 16b). In Fig. 16b, the results are comparable to those obtained for the total integrated intensity (Fig. 16a) but the result is less convincing when we look at a few points (Fig. 9). One should note however that this particular hyperfine transition suffers from noticeable excitation temperature deviations (Caselli et al. 1995) which could explain such a difference on a few data points. In any case, the north-south cut of the ($J_{F_1F_2}:1_{11}-0_{01}$) transition does not reveal the dust peak and in conclusion, optical thickness cannot explain our results. Depletion of N_2 is the only possibility left.

To evaluate the range of densities where N_2H^+ is present, we can try to analyze the N_2H^+ spectra with CLASS HFS method combined with a crude non-LTE (LVG) analysis of the physical conditions, using Green’s collision coefficients (which do not take into account the hyperfine splitting, Green 1975). In the case of the spectrum taken towards the dust peak (Fig. 17), we get a total opacity of ~ 20 (i.e. individual hyperfine component opacities from 0.8 to 5), an excitation temperature of 4.6 K and a remarkably narrow linewidth (0.18 km s^{-1}). Considering a single hyperfine transition as an isolated line, the LVG model indicates a local density of $\sim 2.5 \times 10^4 \text{ cm}^{-3}$ and a total column density of $\sim 10^{13} \text{ cm}^{-2}$ if we suppose the kinetic temperature to be 10 K. If we try to fit the ($J_{F_1F_2}:1_{11}-0_{01}$) transition, we find a density $\sim 8 \times 10^4 \text{ cm}^{-3}$ instead, and towards other places in the cloud, where the N_2H^+ line is not so optically thick, the same crude LVG analysis yields densities around 10^5 cm^{-3} . It thus indicates that towards the main dust core we are mostly measuring the outer parts of the N_2H^+ region where the density is low. This is due to the fact that the escaping photons are those emitted in the last absorbing layer outside the core, the one with the lowest density as the core density decreases radially (Paper II). This density of $2.5 \times 10^4 \text{ cm}^{-3}$ is consistent with the density of $3 \times 10^4 \text{ cm}^{-3}$ we have found for $C^{18}O$ and CS (paper I) together with the fact that CO and N_2H^+ cannot coexist : N_2H^+ appears where CO disappears. Towards the center, it is difficult to evaluate precisely at which density the ion starts to disappear by depletion but the turn-over point at $A_V \approx 50$ mag in Fig. 16 corresponds to a density of $\sim 4 \times 10^5 \text{ cm}^{-3}$ in Fig. 3 of Paper II. Because the dust

⁴ The Grenoble data reduction package. The HFS method assumes LTE holds among the hyperfine components which is approximate in the case of N_2H^+ ($J:1-0$) (Caselli et al. 1995).




Fig. 16. **ab** N_2H^+ ($J:1-0$) integrated intensity pixels cross-correlated with the A_V dust map pixels from Paper II. **a** (left) all seven hyperfine components have been included. **b** (right) only the weakest, optically thin ($J_{F_1F}:1_{11}-0_{01}$) component has been included. A_V map resolution is degraded to $1'$ for KP 12-m data and to $23''$ for IRAM 30-m data

peak maximum density is $2 \times 10^6 \text{ cm}^{-3}$, we can, conservatively, claim that N_2H^+ depletion starts below or close to 10^6 cm^{-3} and probably up to a factor of 2 lower.

4.4.2. N_2D^+ and NH_2D

Another way to check the opacity problem is to look at the rarer isotopomer N_2D^+ . Interestingly enough, the $[\text{N}_2\text{D}^+]/[\text{N}_2\text{H}^+]$ abundance ratio is very high towards molecular peak C, ≈ 0.35 (Tin e et al. 2000). This large deuteration enhancement has been seen for many species in CO depleted cores (including ND_3 in B1 — Lis et al. 2002 — and CD_3OH in IRAS 16293–2422 — Parise et al. 2004) and is clearly linked to the large depletion of heavy species (see e.g. Roberts et al. 2003).

Despite its relatively high abundance, the N_2D^+ high electric dipole moment makes it difficult to populate the $J=2$ rotational level of this species in cold, moderately dense dark cloud cores and thus the ($J:3-2$) transition is probably optically thin or close to optical thinness in the present source. Because the case is not so straightforward for the ($J:2-1$) transition and for the para- NH_2D ($J_{KK'}:1_{11}-1_{01}$) transition, we have selected optically thin hyperfine components of both species to see if the maps were different and if we could find the dust peak. We find that the maps of the selected hyperfine components are identical (within the noise) to the maps presented here (Fig. 8). This can be seen in Fig. 9 where the optically thin component integrated intensities trace the same profile as the total integrated intensities for these two lines. In Fig. 8, we see the molecular peak $30''$ north of the dust peak and a small extension to the dust peak for both NH_2D and N_2D^+ but the dust peak is clearly absent. Thus it does not seem possible to find any species to trace the dust peak except H_2D^+ which has been recently discovered in L1544 (Caselli et al. 2003) and also towards this source (Caselli et al. in prep.) and, with even greater observing

difficulties, D_2H^+ , recently discovered in 16293E (Vastel et al. 2004).

From the molecular peak to the dust peak, the dust column density increases by a factor of ~ 1.5 and the peak density by a factor of ~ 5 following the density profile of Paper II (their Fig. 3). In first approximation, the intensity of the line is proportional to its column density (CD) and to the gas density, $n(\text{H}_2)$, as long as the gas density is below the critical density for the transition we observe. This has been checked with an LVG model for the N_2D^+ ($J:3-2$) line and the range of densities implied here. The intensity of the line is thus proportional to the integral along the line of sight of the product $CD \times n(\text{H}_2)$. This integral increases by a factor ~ 4 from the molecular peak to the dust peak, while the N_2D^+ ($J:3-2$) line integrated intensity is approximately constant. Therefore, the line hyperfine components being close to optical thinness, we must conclude that the line of sight depletion reaches ~ 4 compared to the molecular peak. Consequently, the volume depletion must be at least a factor of ~ 15 for these N-bearing species on average if we suppose that we may have missed a 20% (3σ) intensity increase from molecular peak to dust peak in our observations (we expect a factor 4 increase and we may have only 1.2 , $(4-1)/(1.2-1) = 15$, the -1 is due to the undepleted part in front and behind the core)⁵.

Instead of depletion, one could argue that the molecular peak is peculiar with a strong N_2D^+ enrichment. In fact, Fig. 9 reveals that N_2D^+ and NH_2D do follow the dust profile on both sides of the dust peak and that the molecular peak existence and asymmetry with respect to the dust peak are due in fact to the asymmetric dust profile itself. The molecular peak lies on the northern ridge (Pagani et al. 2003, Paper II) at the closest position to

⁵ A more accurate analysis will be performed in a coming paper when the detailed collisional coefficients (including hyperfine structure) of N_2H^+ will be completely computed (Daniel et al. 2003, Daniel et al. in prep.)

the core before depletion sets in. Because of the elongated ridge with a slow density variation, the region of high density without depletion is large enough to stand out as a molecular peak. In all the other directions away from the dust peak, the density drops much faster and therefore the region of maximum density without depletion is too narrow to be revealed. The column density has somewhat the same effect but to a lesser extent as we have seen above (the density changed by a factor 5 while the column density by a factor 1.5 between molecular peak C and dust peak 1). Therefore, the molecular peak is indeed displaced to the north by $30''$ due to depletion.

4.5. Depletion trigger

Until now, it has not been clear whether the main trigger for depletion of species such as CO and CS is dust opacity (i.e. protection against evaporation by UV radiation) or density (i.e. speed of depletion). It seems from this study that density is the major trigger as the most detailed studies published today quote densities of the order a few 10^4 cm^{-3} very close to our own value of $3 \times 10^4 \text{ cm}^{-3}$ which is the density we have found in Paper I to explain C^{18}O , C^{32}S and C^{34}S line intensities towards the reference position. Tafalla et al. (2002) find $3\text{--}6 \times 10^4 \text{ cm}^{-3}$, Tafalla et al. (2004) find 2.5 and $7.8 \times 10^4 \text{ cm}^{-3}$, and Jørgensen et al. (2004) find $3 \times 10^4 \text{ cm}^{-3}$. On the contrary optical extinction thresholds are usually found close to $A_V = 10$ mag (see Sect. 4.1) while we report $A_V = 20$ mag here (and this is not even sufficient for peak 5—peak E which shows no depletion). Moreover we have recently found in a cloud in the Taurus region C^{18}O depletion starting at $A_V \sim 4$ mag, the lowest A_V value presently known for CO depletion (Boudet et al., in prep.).

Concerning the two important nitrogen-bearing species (N_2H^+ and NH_3 , the latter being represented by its deuterated isotopomer NH_2D only), we find that they are depleted at densities somewhere between 5×10^5 and 10^6 cm^{-3} . While Bergin & Langer (1997) predict such an effect at this density level for grains covered by polar (H_2O) ices, a more recent paper (Aikawa et al. 2003) estimates that N_2H^+ and NH_3 depletion is unlikely below densities of $3 \times 10^7 \text{ cm}^{-3}$. The possible reason which they invoke for this effect is that the binding energy of N_2 onto grains could be lower than usually advocated (a similar effect is found by Bergin & Langer 1997 for grains with no ice or apolar (CO) ice). The present result would tend to show that the polar ice case described by Bergin & Langer (1997) is probably closer to the truth.

4.6. The peak E case

Despite a peak visual extinction of ~ 25 mag (Paper II), Peak E shows no CO depletion and correlatively very weak N_2H^+ emission. This peak is in fact the C^{17}O and C^{18}O emission peak of the whole cloud. This peak is also a 100 and 200 μm peak (Paper I) from which we have derived

an average temperature of ~ 15 K. However, this temperature yields a 200 μm -derived dust extinction $A_V \approx 7$ mag (Paper I) too low compared to the NIR-determined visual extinction (Paper II). This probably indicates that the bulk of the dust is at a lower temperature and that some layers are at higher temperatures than 15 K. These warm layers would not allow depletion of apolar molecules like CO but their total opacity being low, they cannot account for all the C^{18}O emission detected. In Paper I, we have tried to fit the C^{18}O (J:1–0), (J:2–1) and (J:3–2) rotational line emission with a Monte-Carlo code. Although we somewhat failed to reproduce the very low (J:2–1) line emission, we are probably close to the solution, in terms of the column density estimate which is $3.5 \times 10^{15} \text{ cm}^{-2}$. Compared to a peak A_V of ~ 25 mag, this yields an abundance of $\sim 1.4 \times 10^{-7}$, which is the standard value recommended by Frerking et al. (1982). We have only weak C^{32}S (J:2–1) line detections towards peak E in sharp contrast with C^{18}O intensity. Though it is yet difficult to derive a density from a single CS transition, keeping the same $\text{C}^{32}\text{S}/\text{C}^{18}\text{O}$ abundance ratio as measured towards the reference position (i.e. ~ 0.005) we find that C^{32}S (J:2–1) and C^{18}O (J:1–0), (J:2–1) and (J:3–2) are reasonably compatible with a core density of $\sim 10^4 \text{ cm}^{-3}$. Thus, density could again be the explanation for the absence of depletion towards Peak E. Why this peak is less dense and (partially but substantially) warmer than the other peaks in this cloud is not yet clear.

In both Figs. 11 & 13, there is a drop (-30%) of intensity with respect to C^{18}O for the last averaged point (corresponding to the strongest C^{18}O emission). This average point contains only C^{18}O positions in the direction of peak E. As we have seen above, the C^{18}O abundance is normal towards peak E, and we could find the same $\text{C}^{32}\text{S}/\text{C}^{18}\text{O}$ abundance ratio as measured towards the reference position provided that the peak density be no higher than $\sim 10^4 \text{ cm}^{-3}$. Though we cannot do a similar excitation analysis for SO, it is most probable that the intensity drop is also explained by the sole density factor. Thus the peak E is probably not peculiar regarding the species abundances. The question then arises of whether the weak N_2H^+ line observed in that direction is due to chemical destruction as predicted by models or only to the fact that the density is so low that it cannot be observed. The N_2H^+ line we have detected is so weak (~ 0.15 K (4σ)) that only the strongest component ($J_{F_1 F_2}:1_{23}-0_{12}$) emerges from the noise and no CLASS HFS analysis can be done. However, we know the density to be probably no more than 10^4 cm^{-3} and therefore we can use the LVG code to crudely estimate the N_2H^+ column density. With $T_{\text{kin}} = 10$ K, we find a ($J_{F_1 F_2}:1_{23}-0_{12}$) line column density of $3 \times 10^{11} \text{ cm}^{-2}$ which yields a total column density of $\sim 1.2 \times 10^{12} \text{ cm}^{-2}$. Compared to the dust extinction of 25 mag towards peak #5, we find a N_2H^+ abundance of 5×10^{-11} . If we repeat the same analysis for a few N_2H^+ observations towards lines of sight with ~ 40 mag of extinction (to stay away from both the depletion threshold and the threshold of destruction by CO), we find an abundance of

$1.2\text{--}2.1 \times 10^{-10}$, 2 to 4 times higher than towards peak E. Therefore, a low volume density is not the unique reason for the quasi-absence of N_2H^+ emission towards peak E. It has also an abundance lower than in the CO depleted region. Several theoretical and observational studies have shown that N_2H^+ appears in the gas when CO depletes (Bergin & Langer 1997, Bergin et al. 2001, Caselli 2002, Aikawa et al. 2003, Jørgensen et al. 2004). This analysis would appear to again confirm this effect.

5. Conclusions

All species containing at least one atom of the CNO group are depleted in the most inner core of L183. CO, CS and SO are largely depleted but their depletion starts only at a relatively high optical extinction ($A_V \approx 20$ mag) unseen elsewhere. On the contrary, the density ($n \approx 3 \times 10^4 \text{ cm}^{-3}$) for which depletion starts is similar to other findings and this result points towards the fact that density is the main parameter to trigger depletion rather than protection against UV radiation. Depletion effects, reported now in many dark clouds, prevent CO from being an accurate mass tracer for cold dense cores, and forbid the search of such cores via its rarer isotopomers (C^{17}O , C^{18}O). Similarly, CS is not a good density tracer in many cold clouds because it survives in the gas phase only at densities much below its critical density and thus its intensity is mostly proportional to its column density. N-bearing species (especially N_2H^+ and NH_3) are better tracers of dense cores but only up to densities below 10^6 cm^{-3} . Beyond that limit, they also disappear (Bergin et al 2001). Like for L1544 and 16293E, only H_2D^+ (Caselli et al. 2003) and D_2H^+ (Vastel et al. 2004) should be observable in the inner core (Caselli et al., in prep).

Finally, species abundances for chemical models such as those provided by Dickens et al. (2000) for L183 are meaningless averages along the line of sight of species which are not necessarily coexistent spatially. Only models including geometry, depletion and radiation transport can nowadays be usefully compared to the detailed observations allowed by modern telescopes. Such models have recently been proposed by Bergin et al. (2001, including a chemical network scheme) or by Jørgensen et al. (2004) and similar studies in protostellar cores have been presented by Tafalla et al. (2002, 2004).

Acknowledgements. J.R. Pardo's research work is supported by Spanish MCyT grants ESP 2003-01627 and AYA 2002-10113-E. We thank the referee, E.D. (Ted) Bergin, for useful comments and suggested improvements to the paper.

References

Aikawa Y., Ohashi N., Herbst E., 2003, ApJ 593, 906
 Alves J., Lada C.J., Lada E.A., 1999, ApJ 515, 265
 Bacmann, A., Lefloch, B., Ceccarelli, C., et al. 2003, ApJL 585, L55
 Bergin E.A., Langer W.D., 1997, ApJ 486, 316

Bergin E.A., Ciardi D.R., Lada, C.J., Alves, J., Lada, E.A., 2001, ApJ 557, 209
 Bergin E.A., Alves, J., Huard, T., Lada, C.J., 2002, ApJ 570, L101
 Caselli, P., Myers, P.C., Thaddeus, P., 1995, ApJL 455, L77
 Caselli P., 2002, P&SS 50, 1133
 Caselli, P., van der Tak, F., Ceccarelli, C., Bacmann, A., 2003, A&A, 403, L37
 Daniel F., Dubernet M.-L., Pagani L., in SF2A Scientific Highlights 2003, eds Combes F., Barret D., Contini T., Pagani L., EdP-Science, Les Ulis, p.205
 Dickens J.E., Irvine W.M., Snell R.L., et al., 2000, ApJ 542, 870
 Frerking, M.A., Langer, W.D., Wilson, R.W., 1982, ApJ 262, 590
 Goldsmith, P. & Langer, W.D., 1999, ApJ 517, 219
 Green, S., 1975, ApJ 201, 366
 Green, S., 1994, ApJ 434, 188
 Jørgensen J.K., Schöier F.L., van Dishoeck E.F., 2004, A&A, in press
 Juvela, M., Mattila, K., Lehtinen, K., et al., 2002, A&A 382, 583
 Kramer, C., Alves, J., Lada, C.J., et al., 1999, A&A 342, 257
 Kutner, M.L., Ulich, B.L., 1981, ApJ 250, 341
 Linke, R. A., Goldsmith, P. F., Wannier, P. G., Wilson, R. W., & Penzias, A. A., 1977, ApJ 214, 50
 Lis, D.C., Roueff, E., Gerin, M., et al., 2002, ApJ 571, L55
 Pagani, L., Gallego, A.T., Apponi, A.J., 2001, A&A 380, 384 (+erratum)
 Pagani, L., Pardo, J.-R., Fich, M., Motte, F., and Stepnik, B., 2002, in "Infrared and Submillimeter Space Astronomy", EAS Series, Vol. 4, Giard M. et al., eds, held in Toulouse, June 2001, EdP Sciences, Les Ulis, p.145 (Paper I)
 Pagani, L., Lagache, G., Bacmann, A., et al. 2003, A&A 406, L59
 Pagani, L., Bacmann, A., Motte, F., et al. 2004, A&A 417, 605 (Paper II)
 Penzias A.A., 1975, In: Atomic and molecular physics and the interstellar matter; Proceedings of the Twenty-sixth Summer School of Theoretical Physics, Les Houches, France, July 1-August 23, 1974. Amsterdam, North-Holland Publishing Co., 1975, p. 373-408.
 Parise, B., Castets, A., Herbst, E., et al. 2004, A&A 416, 159
 Roberts, H., Herbst, E., Millar, T.J., 2003, ApJ 591, L41
 Swade, D.A., 1989a, ApJS, 71, 219
 Swade, D.A., 1989b, ApJ, 345, 828
 Tafalla, M., Myers, P.C., Caselli, P., Walmsley, C.M., Comito, C., 2002, ApJ 569, 815
 Tafalla, M., Myers, P.C., Caselli, P., Walmsley, C.M., 2004, A&A in press
 Teyssier, D., Hennebelle, P., Pérault, M., 2002, A&A, 382, 624
 Tiné, S., Roueff, E., Falgarone, E., Gerin, M., Pineau des Forêts, G., 2000, A&A 356, 1039
 Vastel, C., Phillips, T. G., Yoshida, H., 2004, ApJ 606, L127
 Willacy K., Langer W.D., Velusamy T., 1998, ApJ 507, L171

Appendix A: Subthermally excited optically thick lines

A radiotelescope which observes a source with opacity τ and excitation temperature T_{ex} by comparison with an empty field, detects an antenna temperature T_a increase given by




Fig. A.1. $C^{32}S$ (J:1-0) and (J:2-1) transition properties as a function of column density. LVG model with $T_{\text{kin}} = 10$ K, $\delta v = 1$ km s $^{-1}$ and $n = 10^3$ cm $^{-3}$




Fig. A.2. Same as Fig. A.1 except $T_{\text{kin}} = 50$ K

$$\Delta T_a = \eta(1 - e^{-\tau})(J(T_{\text{ex}}) - J(T_{\text{bg}})) \quad (\text{A.1})$$

with η the coupling efficiency of the telescope, T_{bg} , the temperature of the cosmological background radiation and

$J(T)$ is given by

$$J(T) = \frac{h\nu}{k} \frac{1}{e^{\frac{h\nu}{kT}} - 1} \quad (\text{A.2})$$

where h is the Planck constant, k the Boltzmann constant and ν the line frequency. When $\tau \gg 1$, eq. A.1 simplifies to

$$\Delta T_a = \eta(J(T_{\text{ex}}) - J(T_{\text{bg}})) \quad (\text{A.3})$$

which would seem to be independent of τ and therefore of the species column density as τ is defined by

$$\tau = \frac{c^2 g_u A_{ul} N_l}{8\pi g_l \nu^2 \Delta\nu} \left[1 - e^{-\frac{h\nu}{kT_{\text{ex}}}} \right] \quad (\text{A.4})$$

with c the light speed, A_{ul} the Einstein spontaneous emission coefficient, g_u and g_l the upper and lower species level statistical weights and N_l , the lower level column density. The latter is related to the total column density via the partition function.

In fact, when a line is optically thick and the local density is much below the critical density (defined by $n_{\text{crit}} = A_{ul}/\gamma_{ul}$, γ_{ul} being the collision probability), each exciting collision has a higher probability to be followed by a radiative decay rather than a collisional deexcitation. But similarly, if the photon is reabsorbed because of high opacity, it will still be reemitted rather than having the species be deexcited by a collision. Thus each exciting collision will give birth to a photon which will eventually escape. If the opacity is high, many photons are retained in the cloud before escaping and the net result is an increase of the excitation temperature and therefore of the antenna temperature in eq. A.3. We refer the reader to Penzias (1975) or Linke et al. (1977) for a mathematical demonstration. Here we want to illustrate this result via two simulations and actual observations.

Figs A.1 and A.2 show the case of $C^{32}S$ (J:1-0) and (J:2-1) lines in a very low density medium (10^3 cm $^{-3}$) and at two different kinetic temperatures (Fig. A.1, $T_{\text{kin}} = 10$ K, Fig. A.2, $T_{\text{kin}} = 50$ K). The line opacity and antenna temperatures are proportional to the total column density below $N = 10^{13}$ cm $^{-2}$, which corresponds to $\tau \lesssim 1$ and T_{ex} is almost constant. Above $N = 10^{13}$ cm $^{-2}$, i.e. $\tau > 1$, the excitation temperature starts to rise and the antenna temperature keeps increasing linearly first then deviates while approaching the excitation limit (collisional deexcitation starts to be efficient when too many molecules are in the excited state).

In L183, the N_2H^+ (J:1-0) line has the advantage of having both a hyperfine transition always optically thin ($J_{F_1 F_2}:1_{11}-0_{01}$) and another one, almost always optically thick ($J_{F_1 F_2}:1_{23}-0_{12}$) in the range of conditions offered by this cloud. The line has a critical density around 10^6 cm $^{-3}$ and is therefore subthermally excited everywhere in the cloud. Fig. A.3 shows the comparison of the two hyperfine transition integrated intensities together with the 1:7 slope corresponding to their LTE ratio. The proportionality is clearly visible despite a slight departure from the 1:7




Fig. A.3. Comparison of the N_2H^+ ($J_{F_1 F_2}:1_{10}-0_{01}$) and ($J_{F_1 F_2}:1_{23}-0_{12}$) line integrated intensities. Square symbols identify those spectra for which the ($J_{F_1 F_2}:1_{23}-0_{12}$) line is optically thin. Low signal-to-noise ratios explain the presence of “optically thick” crosses in the bottom left corner

slope. This departure is probably due to the emergence of a saturation effect or to the fact that the ($J_{F_1 F_2}:1_{11}-0_{01}$) line often departs from the LTE ratio (Caselli et al. 1995).